



Chronic Spontaneous Urticaria (CSU)
with hives not controlled by H1 antihistamine

DUPIXENT[®] 
(dupilumab) Injection
200mg • 300mg



Help with Chronic
Spontaneous Urticaria?



First and only FDA-approved biologic
for appropriate patients as young as 2 years
old with chronic spontaneous urticaria.

INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 2 years of age and older with chronic spontaneous urticaria (CSU) who continue to have hives that are not controlled with H1 antihistamine treatment. **DUPIXENT is not used** to treat any other forms of hives (urticaria). It is not known if DUXIPENT is safe and effective in children with CSU under 2 years of age, or who weigh less than 11 pounds (5 kg).

**Please see Important Safety Information
throughout and accompanying full Prescribing
Information including Patient Information.**

What is CSU?

Chronic Spontaneous Urticaria (CSU),

also known as chronic hives with no known cause, may happen when the immune system is out of balance.

Key characteristics include:

Itchy red or skin-colored bumps with no known cause

6+ weeks of hives on and off, with or without swelling deep in the skin

Spontaneous hives that appear then disappear, often on the chest, back, arms, or legs

No known trigger

With CSU, inflammation under the skin contributes to spontaneous flare-ups of itch and hives

that are not related to an allergy or external trigger.

IMPORTANT SAFETY INFORMATION:

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DID YOU KNOW?

~50%

of adults find antihistamines are not enough

2-3 years

is the average time adults spend seeking a diagnosis

2-5 years

is the average length of most adult cases, but can last longer

Q Why are H1 antihistamines sometimes not enough?

You may have been taking antihistamines steadily, and for a long time.

Maybe you're even taking more than the standard dose, and are still stuck in a cycle of flares of itch and hives, antihistamines, then more flares.

However, antihistamines may not be enough to control your CSU symptoms.

“

The itch was absolutely unbearable.

I would just cry because I itched so bad. It never really went away and just felt like it was dominating my life.

—Emily A., actual DUPIXENT patient”

IMPORTANT SAFETY INFORMATION:

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

Q What is DUPIXENT?

DUPIXENT IS AN INJECTABLE MEDICINE THAT TARGETS A KEY SOURCE OF INFLAMMATION THAT CAN CONTRIBUTE TO CSU

- DUPIXENT continuously works within your body between doses to help manage CSU symptoms.
- It can help you get ahead of your unpredictable symptoms by reducing the intense itch and number of hives.

DUPIXENT IS NOT:

- ✗ A steroid
- ✗ A daily pill
- ✗ An immunosuppressant

IMPORTANT SAFETY INFORMATION:

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

DUPIXENT[®]
(dupilumab) Injection
200mg • 300mg



IMPORTANT SAFETY INFORMATION:

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have CSU and asthma and use an asthma medicine. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

What kind of results can I expect?

DUPIXENT was studied in two 24-week clinical trials in 284 patients ages 12 years and older with CSU who continued to have hives despite taking H1 antihistamines. All patients remained on H1 antihistamines during the trials.



AT 24 WEEKS:

LESS ITCH*

Trial 1:

64% itch reduction with DUXIPENT
vs 35% on placebo

Trial 2:

48% itch reduction with DUXIPENT
vs 33% on placebo

REDUCTION IN BOTH ITCH AND HIVES†

Trial 1:

66% reduction
on DUXIPENT
vs 36% on placebo

Trial 2:

48% reduction
on DUXIPENT
vs 32% on placebo

DUPIXENT's use for treating CSU in children ages 2-11 years is supported by the results from two studies that included patients 12 years of age and older with CSU.

*Patients rated itch daily (from 0-3) for 7 days (max score 21), compared to baseline score.

†Patients also rated their hives daily (from 0-3) for 7 days (max score 21). This score was added to their itch severity score, compared to baseline score.

DUPIXENT®
(dupilumab) Injection
200mg • 300mg

Nearly twice as many patients had no itch and no hives at 24 weeks

Trial 1:

32% on DUXIPENT
vs 13% on placebo

Trial 2:

30% on DUXIPENT
vs 17% on placebo



AS EARLY AS 3 WEEKS:

Reduced itch and hives as early
as week 3 in some patients

In both trials, patients saw 32% reduction with DUXIPENT vs 23% with placebo.

Results at Week 3 were evaluated after the clinical trials, and any results at Week 3 may be due to chance.

Individual results may vary.

The most common side effects include injection site reactions.

IMPORTANT SAFETY INFORMATION:

DUPIXENT can cause serious side effects, including:

- **Allergic reactions.** DUXIPENT can cause allergic reactions, including skin reactions, that can sometimes be severe. Stop using DUXIPENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DUPIXENT[®]

(dupilumab) Injection
200mg • 300mg

Q How do I take DUPIXENT?

DUPIXENT IS AVAILABLE AS A PRE-FILLED PEN OR PRE-FILLED SYRINGE



- » DUPIXENT is a biologic medicine given under the skin (subcutaneous injection)
- » It is not taken every day. DUPIXENT is taken only once or twice a month (based on age and weight)
 - Your doctor will tell you how much DUPIXENT to take
- » DUPIXENT can be given by your doctor or self-administered at home
 - Do not try to prepare and inject DUPIXENT until being properly trained by your doctor
 - In children 12 years of age and older, it is recommended that DUPIXENT be given by or under the supervision of an adult
 - In children 2 years to less than 12 years of age, DUPIXENT should be given by a caregiver



Click here to
learn more about how
to inject DUPIXENT.

Your specialist will prescribe the appropriate dosage based on the recommendations below:

ADULTS (18+ YEARS OF AGE)

BODY WEIGHT	INITIAL DOSE	SUBSEQUENT DOSAGE
Any weight	600 mg (two 300-mg injections)*	300 mg every 2 weeks

CHILDREN AND TEENS (6-17 YEARS OF AGE)

BODY WEIGHT	INITIAL DOSE	SUBSEQUENT DOSAGE
132 lb or more	600 mg (two 300-mg injections)*	300 mg every 2 weeks
66 lb to under 132 lb	400 mg (two 200-mg injections)*	200 mg every 2 weeks
33 lb to under 66 lb	600 mg (two 300-mg injections)*	300 mg every 4 weeks

*Administered into different injection sites.

YOUNG CHILDREN (2-5 YEARS OF AGE)

BODY WEIGHT	DOSAGE [†]
33 lb to under 66 lb	300 mg (one 300-mg injection) every 4 weeks
11 lb to under 33 lb	200 mg (one 200-mg injection) every 4 weeks

[†] For pediatric patients 2 to 5 years of age with CSU, no initial loading dose is recommended.

IMPORTANT SAFETY INFORMATION:

DUPIXENT can cause serious side effects, including (cont'd):

- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

How can **DUPIXENT** MyWay® support me?

**PAY AS
LITTLE
AS \$0***



The **DUPIXENT** MyWay Copay Card may help reduce the out-of-pocket cost of **DUPIXENT** for eligible patients. If you have commercial health insurance, you may be eligible to pay as little as \$0* in copay per fill of **DUPIXENT** through the **DUPIXENT** MyWay Copay Card.

Terms and conditions apply.



**CLICK HERE TO
LEARN MORE ABOUT
DUPIXENT MyWay**

Or call to enroll: 1-844-DUPIXENT
(1-844-387-4936), Option 1

*Subject to the program maximum per patient per calendar year.

Approval is not guaranteed. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. **DUPIXENT** MyWay reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford **DUPIXENT**.

Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.



MAY BE ELIGIBLE FOR \$0* COPAY IF YOU:

- » Have commercial insurance
- » Have a **DUPIXENT** prescription for an FDA-approved condition
- » Are a resident of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the U.S. Virgin Islands; and
- » Are a patient or caregiver aged 18 years or older

INELIGIBLE IF YOU:

Have government insurance—Medicaid, Medicare, Veterans Affairs, Department of Defense, TRICARE, or other federal or state programs. **Ineligible?** **DUPIXENT** MyWay may be able to provide alternative financial assistance for eligible patients.

IMPORTANT SAFETY INFORMATION:

The most common side effects in patients with chronic spontaneous urticaria include injection site reactions.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of **DUPIXENT**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.




DUPIXENT MyWay[®] can help support you throughout your treatment journey

The DUPIXENT MyWay Patient Support Program is designed to help you get DUPIXENT[®] (dupilumab) as quickly as possible and stay on track with therapy.

You can obtain information about your insurance coverage for DUPIXENT and sign up for:

- **Supplemental injection support and training** with Nurse Educators in person, virtually, or by phone
- **Injection and prescription reminders** that can help you remember to take your next dose and refill your prescription
- **Insightful tips, tools, and resources** designed to help you along your journey



**CLICK HERE TO
STAY ON TRACK WITH
THE *DUPIXENT MyWay* APP**

IMPORTANT SAFETY INFORMATION:

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 2 years to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

IF H1 ANTIHISTAMINES AREN'T GIVING
YOU THE CONTROL YOU NEED,
ASK YOUR DERMATOLOGIST OR
ALLERGIST ABOUT

DUPIXENT[®] 
(dupilumab) Injection
200mg • 300mg



[Click here](#) for help with cost,
coverage, and your next
steps with DUPIXENT.

IMPORTANT SAFETY INFORMATION:

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

sanofi | **REGENERON**[®]